



## ANALYTICAL REPORT

<b>Client</b>	Blackcurrants New Zealand Ltd
<b>Sample(s) Analysed</b>	Five frozen berry samples were analysed for individual anthocyanins, total anthocyanins, and vitamin C content by HPLC and for antioxidant capacity using the ORAC <sub>FL</sub> , FRAP and Folin-Ciocalteu assays.
<b>Date Received</b>	11 July 2007
<b>HortResearch Ref</b>	EA56
<b>Results</b>	Refer to page 2 for result tables
<b>Comments</b>	<p>ORAC<sub>FL</sub> Method: Based on Ou et al (Journal of Agriculture and Food Chemistry 49: 4691-4626 2001) an improvement on the method developed by Cao et al (Clinical Chemistry 41: 1738-1744 1995). FRAP Method: Based on protocol used by Scottish Crops Research Institute (Deighton, et al., 2000) as adapted from Benzie and Strain (1996). Folin-Ciocalteu Method: Based on protocol of Coseteng and Lee (1987). Anthocyanin and Vitamin C concentrations were determined by reversed-phase HPLC.</p> <p>Standards: Trolox (water soluble vitamin E analogue); gallic acid; cyanidin-3-glucoside; ascorbic acid</p> <p>FRAP &amp; ORAC reported as mmol of Trolox equivalents per 100g of fruit. Folin-Ciocalteu reported as mmol of gallic acid equivalents per 100g of fruit. Total vitamin C (combined ascorbic acid and dehydroascorbic acid) reported as mg per 100g of fruit. Total anthocyanins reported as mg of cyanidin-3-glucoside equivalents per 100g of fruit. Individual anthocyanins reported as µg of cyanidin-3-glucoside equivalents per gram of fruit.</p>
<b>Analyst</b>	Dan Comeskey
<b>Checked</b>	Tony McGhie
<b>Date</b>	Tuesday, 17 July 2007

### Disclaimer

*HortResearch does not give any warranty or assurance in relation to the accuracy of or fitness for any particular use or application of any result contained in this report. Neither HortResearch, nor any of its employees shall be liable for any cost, claim, liability or the like which may be suffered as a result of reliance on any information contained in this report*

	NZ Blackcurrant Ben Rua	NZ Blackcurrant Ben Ard	Swedish Bilberry	European Bilberry	Canadian Blueberry
FRAP mmol TE/100g	5.22	5.55	3.75	2.85	1.35
ORAC mmol TE/100g	9.51	12.15	12.04	10.20	6.00
Folin-Ciocalteu mmol GAE/100g	660	688	617	502	271
Total Vitamin C mg/100g	191	175	1.4	0.7	5.1
Total Anthocyanins mg/100g	517	584	526	431	135

Anthocyanin µg/g	NZ Blackcurrant Ben Rua	NZ Blackcurrant Ben Ard	Swedish Bilberry	European Bilberry	Canadian Blueberry
Delphindin-galactoside	n.d.	n.d.	685	482	132
Delphindin-glucoside	369	899	688	460	84
Delphinidin-rutinoside	2325	2049	n.d.	n.d.	n.d.
Cyanidin-galactoside + Delphindin-arabinoside	n.d.	n.d.	1163	894	n.d.
Delphindin-arabinoside	n.d.	n.d.	n.d.	n.d.	155
Cyanidin-glucoside	153	456	557	500	12
Petunidin-galactoside	n.d.	n.d.	210	179	75
Cyanidin-rutinoside	2120	2252	n.d.	n.d.	n.d.
Cyanidin-arabinoside	n.d.	n.d.	417	362	14
Petunidin-glucoside	n.d.	n.d.	417	322	78
Petunidin-arabinoside	n.d.	n.d.	207	183	66
Malvidin-galactoside	n.d.	n.d.	390	407	220
Malvidin-glucoside	n.d.	n.d.	422	396	163
Malvidin-arabinoside	n.d.	n.d.	85	94	197
Minor Anthocyanins	208	180	16	29	151
Total Anthocyanins	5175	5836	5257	4310	1348